arXiv:1512.00049v1 [math.AC] 30 Nov 2015

A NOTE ON THE GROWTH OF REGULARITY WITH RESPECT TO FROBENIUS

WENLIANG ZHANG

Abstract. Let $R = k[x_1, \dots, x_n]/I$ be a graded k-algebra where k is a field of prime characteristic and let J be a homogeneous ideal in R. Denote $(x_1, \dots, x_n)$ by $\mathfrak{m}$. We prove that there is a constant C (independent of e) such that the regularity of $H^s_{\mathfrak{m}}(R/J^{[p^e]})$ is bounded above by Cp^e for all $e \geq 1$ and all integers s such that $s+1$ is at least the dimension of the locus where R/J doesn't have finite projective dimension.

1. introduction

Let $S = k[x_1, \dots, x_n]$ be a polynomial ring over a field k of characteristic $p > 0$ with the standard grading. Let $\mathfrak{m}$ be the irrelevant maximal homogeneous ideal of S. For each graded artinian S-module N, we set $a(N) := \max\{j \mid N_j \neq 0\}$. And the Castelnuovo-Mumford regularity of a finitely generated graded S-module M is defined as

$$\operatorname{reg}(M) := \max\{a(H^j_{\mathfrak{m}}(M)) + j \mid j \in \mathbb{Z}\}.$$

For each ideal $\mathfrak{a}$ of S, let $\mathfrak{a}^{[p^e]}$ denotes the ideal generated by $\{r^{p^e} \mid r \in \mathfrak{a}\}$. [Kat98] asked the following question:

Question 1.1. Let I, J be graded ideals of S. Does there exist a constant C such that $\operatorname{reg}(\frac{S}{I+J^{[p^e]}}) \leq Cp^e$ for all $e \geq 1$?

When such a constant exists, we say that $\operatorname{reg}(\frac{S}{I+J^{[p^e]}})$ grows linearly with respect to p^e (or with respect to Frobenius). While the original motivation behind Question 1.1 concerns whether tight closure commutes with localization at a single element, it was discovered in [KZ14] that a positive answer to Question 1.1 will imply that the F-jumping coefficients of any ideal in a finitely generated k-algebra are discrete. A positive answer to Question 1.1 is known in some cases; for instance, when $\dim(\operatorname{Sing}(S/I) \cap V(J)) \leq 1$ ([Cha07], [Bre05], [KZ14]) and when S/I has finite graded Frobenius representation type ([KSSZ14]).

Let $R = k[x_1, \dots, x_n]/I$ be a standard graded ring over a field k of characteristic $p > 0$ and $\mathfrak{m} = (x_1, \dots, x_n)$. Let $F : R \longrightarrow R$ denote the Frobenius endomorphism and $F^e_* R$ denote the target of F^e. For each $e \geq 1$, the R-module $F^e_* R$ is graded via

$$\deg(F^e_* r) = \frac{1}{p^e} \deg(r)$$

for every homogeneous element $r \in R$. Consequently, for each graded R-module M, the R-module $M \otimes_R F^e_* R$ is graded by $\deg(z \otimes F^e_* r) = \deg(z) + \frac{1}{p^e} \deg(r)$.

Our main result is the following theorem:

Theorem 1.2. *Let $R = k[x_1, \dots, x_n]/I$ be a standard graded ring over a field k of characteristic $p > 0$ and M be a finitely generated graded R=module. Set*

$$Npd(M) = \{\mathfrak{p} \in \operatorname{Spec}(R) \mid M_{\mathfrak{p}} \text{ does not have finite projective dimension over } R_{\mathfrak{p}}\}.$$

Then there is a constant C such that

$$a(H^s_{\mathfrak{m}}(M \otimes_R F^e_* R)) \leq C$$

for all $s \geq \dim(Npd(M)) - 1$ and all $e \geq 1$.

The author is partially supported by the National Science Foundation through grant DMS #1405602.

An immediate corollary is:

Corollary 1.3. *Let $R = k[x_1, \ldots, x_n]/I$ be a standard graded ring over a field k of characteristic $p > 0$ and J be a homogeneous ideal of R. Then $a(H^s_{\mathfrak{m}}(R/J^{[p^e]}))$ grows linearly with respect to p^e for each $s \geq \dim(Npd(R/J)) - 1$.*

In particular, $a(H^s_{\mathfrak{m}}(R/J^{[p^e]}))$ grows linearly with respect to p^e for $s \geq \dim(R/J) - 1$.

ACKNOWLEDGEMENT

We would like to thank Mordechai Katzman for inspiring discussions.

2. Proof of Theorem 1.2

Remark 2.1. It is well known that there is a degree-preserving isomorphism $R_f \otimes_R F^e_* R \xrightarrow{a \otimes r' \mapsto a^{p^e} r'} F^e_* R_f$ for each homogeneous element $f \in R$. Consequently, we have

$$H^j_{\mathfrak{m}}(F^e_* R) \cong F^e_* H^j_{\mathfrak{m}}(R).$$

Let $a(H^j_{\mathfrak{m}}(F^e_* R))$ denote $\max\{\rho \in \mathbb{Q} \mid H^j_{\mathfrak{m}}(F^e_* R)_\rho \neq 0\}$. It follows immediately from Remark 2.1 that

Proposition 2.2. $a(H^j_{\mathfrak{m}}(F^e_* R)) = \frac{1}{p^e} a(H^j_{\mathfrak{m}}(R))$.

Lemma 2.3. *For each finitely generated graded R-module M, there is a constant C_{ij} (independent of e) such that $a(\operatorname{Tor}^i_R(H^j_{\mathfrak{m}}(F^e_* R), M)) \leq C_{ij}$ for all e.*

Proof. Let $\mathcal{F}_\bullet$ be a graded free resolution of M. Write $\mathcal{F}_i = \oplus_{n_i} R(\alpha_{n_i})$. Then

$$\mathcal{F}_i \otimes_R H^j_{\mathfrak{m}}(F^e_* R) \cong \oplus_{n_i} H^j_{\mathfrak{m}}(F^e_* R)(\alpha_{n_i}).$$

The conclusion now follows from Proposition 2.2. □

Consider the complex $\mathcal{C}_\bullet$ which is the Čech complex associated with $x_1, \ldots, x_n$ with the indices reversed as follows:

$$0 \leftarrow \mathcal{C}_0 \leftarrow \mathcal{C}_1 \leftarrow \cdots \leftarrow \mathcal{C}_n \leftarrow 0$$

where $\mathcal{C}_j = \check{C}_{n-j}(\underline{x}; R)$. Hence $H^j(\mathcal{C}_\bullet \otimes_R N) = H^{n-j}_{\mathfrak{m}}(N)$ for each R-module N.

Let M be a finitely generated graded R-module and $\mathcal{F}_\bullet$ be a graded free resolution of M. Consider the double complex $\mathcal{C}_{\bullet,\bullet}$:


0 ← $\mathcal{F}_2 \otimes_R \mathcal{C}_0 \otimes_R F^e_* R$ ← $\mathcal{F}_2 \otimes_R \mathcal{C}_1 \otimes_R F^e_* R$ ← ⋯
0 ← $\mathcal{F}_1 \otimes_R \mathcal{C}_0 \otimes_R F^e_* R$ ← $\mathcal{F}_1 \otimes_R \mathcal{C}_1 \otimes_R F^e_* R$ ← ⋯
0 ← $\mathcal{F}_0 \otimes_R \mathcal{C}_0 \otimes_R F^e_* R$ ← $\mathcal{F}_0 \otimes_R \mathcal{C}_1 \otimes_R F^e_* R$ ← ⋯
0
0

Note that $\mathcal{C}_{\bullet,\bullet}$ depends on e.

Lemma 2.4. *For each integer α, there is a constant C_α (independent of e) such*

$$a(H_\alpha(\operatorname{Tot}(\mathcal{C}_{\bullet,\bullet}))) \leq C_\alpha.$$

Proof. It is clear that $\mathcal{C}_{\bullet,\bullet}$ is a first quadrant double complex and hence it induces two spectral sequences both of which converge to the homology of the total complex ([Wei94, § 5.6]). One of these spectral sequence comes from taking homology horizontally first and then vertically, whose E^2-page is given by $\operatorname{Tor}_i^R(H^j_{\mathfrak{m}}(F^e_*R), M)$. Since $H^j_{\mathfrak{m}}(F^e_*R) = 0$ for all $j \geq d+1$ with $d = \dim(R)$, the E^∞-page is the same as the E_{d+1}-page. Hence there are only finitely many $\operatorname{Tor}_i^R(H^j_{\mathfrak{m}}(F^e_*R), M)$ contributing to $H_\alpha(\operatorname{Tot}(\mathcal{C}_{\bullet,\bullet}))$ for each integer α. Lemma 2.3 finishes the proof. □

To prove our main result, we need the following theorem due to Peskine-Szpiro [PS73]:

Theorem 2.5 (Peskine-Szpiro Acyclicity Lemma)**.** *Let R be a noetherian commutative ring of characteristic $p > 0$ and N be a finite R-module. If N has finite projective dimension, then*

$$\operatorname{Tor}_i^R(F^e_*R, N) = 0$$

for all $i \geq 1$ and all e.

Recall that $Npd(M)$ denote the locus where M does not have finite projective dimension, *i.e.*

$$Npd(M) = \{\mathfrak{p} \in \operatorname{Spec}(R) \mid M_{\mathfrak{p}} \text{ does not have finite projective dimension over } R_{\mathfrak{p}}\}.$$

Proof of Theorem 1.2. We consider the other spectral sequence induced by the double complex $\mathcal{C}_{\bullet,\bullet}$, *i.e.* by taking homology vertically first and then horizontally:

$$E^2_{i,j} = H^{n-i}_{\mathfrak{m}}(\operatorname{Tor}_j^R(F^e_*R, M)) \Rightarrow H_{i+j}(\operatorname{Tot}(\mathcal{C}_{\bullet,\bullet})).$$

The differential coming into $H^s_{\mathfrak{m}}(M \otimes_R F^e_*R) = E^2_{n-s,0}$ is 0 since $E^2_{n-s+2,-1} = 0$. Set $\dim(Npd(M)) = t$. We can see that $\dim(\operatorname{Tor}_1(F^e_*R, M)) \leq t$ by Theorem 2.5. Hence, if $s \geq t-1$, we have $H^{s+2}_{\mathfrak{m}}(\operatorname{Tor}_1(F^e_*R, M)) = 0$ which implies that the differential going out of $H^s_{\mathfrak{m}}(M \otimes_R F^e_*R) = E^2_{n-s,0}$ is also 0. Therefore, if $s \geq t-1$, we have

$$H^s_{\mathfrak{m}}(M \otimes_R F^e_*R) = E^\infty_{n-s,0}.$$

According to Lemma 2.4, there is a constant C_{n-s} such that $a(H_{n-s}(\operatorname{Tot}(\mathcal{C}_{\bullet,\bullet}))) \leq C_{n-s}$. Since $E^\infty_{n-s,0}$ is a subquotient of $H_{n-s}(\operatorname{Tot}(\mathcal{C}_{\bullet,\bullet}))$, one can see that

$$a(H^s_{\mathfrak{m}}(M \otimes_R F^e_*R)) \leq C_{n-s}.$$

This finishes the proof. □

Proof of Corollary 1.3. It is well known that there is a degree-preserving isomorphism of F^e_*R-modules (hence R-modules):

$$R/J \otimes_R F^e_*R \xrightarrow{\bar{r}\otimes r' \mapsto \overline{r^{p^e} r'}} F^e_*R/J^{[p^e]}$$

for each graded ideal J of R, where the grading on F^e_*M is given by $\deg(F^e_* z) = \frac{1}{p^e}\deg(z)$ for any graded R-module M. It follows from Theorem 1.2 that there is a constant C such that

$$a(H^s_{\mathfrak{m}}(R/J \otimes_R F^e_*R)) \leq C$$

for each $s \geq \dim(Npd(R/J)) - 1$ and all $e \geq 1$. Therefore,

$$a(H^s_{\mathfrak{m}}(R/J^{[p^e]})) = p^e a(F^e_* H^s_{\mathfrak{m}}(R/J^{[p^e]})) = p^e a(H^s_{\mathfrak{m}}(F^e_*R/J^{[p^e]}) = p^e a(H^s_{\mathfrak{m}}(R/J \otimes_R F^e_*R)) \leq Cp^e$$

for each $s \geq \dim(Npd(R/J)) - 1$ and all $e \geq 1$. □

Remark 2.6. Since $Npd(R/J) \subset \operatorname{Sing}(R) \cap \operatorname{Supp}(R/J)$, Corollary 1.3 recovers [KZ14, 3.5].

References

[Bre05] H. Brenner: *A linear bound for Frobenius powers and an inclusion bound for tight closure*, Michigan Math. J. **53** (2005), no. 3, 585–596. 2207210 (2006k:13010)

[Cha07] M. Chardin: *On the behavior of castelnuovo-mumford regularity with respect to some functors*, arXiv:0706.2731.

[Kat98] M. Katzman: *The complexity of Frobenius powers of ideals*, J. Algebra **203** (1998), no. 1, 211–225. 1620654 (99e:13003)

[KSSZ14] M. Katzman, K. Schwede, A. K. Singh, and W. Zhang: *Rings of Frobenius operators*, Math. Proc. Cambridge Philos. Soc. **157** (2014), no. 1, 151–167. 3211813

[KZ14] M. Katzman and W. Zhang: *Castelnuovo-Mumford regularity and the discreteness of F-jumping coefficients in graded rings*, Trans. Amer. Math. Soc. **366** (2014), no. 7, 3519–3533. 3192605

[PS73] C. Peskine and L. Szpiro: *Dimension projective finie et cohomologie locale. Applications à la démonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck*, Inst. Hautes Études Sci. Publ. Math. (1973), no. 42, 47–119. 0374130 (51 #10330)

[Wei94] C. A. Weibel: *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. 1269324 (95f:18001)

Department of Mathematics, Statistics, and Computer Science, University of Illinois, Chicago, IL 60607, USA

E-mail address: wlzhang@uic.edu